


Diverticular Disease



Sharon R. Roseman, MD

Practice Limited to Gastroenterology

701 Broad Street, Suite 411 ♦ Sewickley, PA 15143

PH: 412-749-7160 ♦ FAX: 412-749-7388

What Is Diverticulosis?

Most people have in their colons small pouches that bulge outward through weak spots, like an inner tube that pokes through weak places in a tire. Each pouch is called a diverticulum. The condition of having diverticula is called diverticulosis. About half of all Americans age 60 to 80, and almost everyone over age of 80, have diverticulosis.

When the pouches become infected or inflamed, the condition is called diverticulitis. This happens in 10 to 25 percent of people with diverticulosis. Diverticulosis and diverticulitis are also called diverticular disease.

What Causes Diverticular Disease?

Doctors believe a low-fiber diet is the main cause of diverticular disease. The disease was first noticed in the United States in the early 1900's. At about the same time, processed foods were introduced to the American diet. Many processed foods contain refined, low-fiber flour. Unlike whole-wheat flour, refined flour has no wheat bran.

Diverticular disease is common in developed or industrialized countries-particularly the United States, England, and Australia-where low-fiber diets are common. The disease is rare in countries of Asia and Africa, where people eat high-fiber vegetable diets.

Fiber is the part of fruits, vegetables, and grains that the body cannot digest. Some fiber dissolves easily in water (soluble fiber). It takes on a soft, jelly-like texture in the intestines. Some fiber passes almost unchanged through the intestines (insoluble fiber). Both kinds of fiber help make stools soft and easy to pass. Fiber also prevents constipation.

Constipation makes the muscles strain to move stool that is too hard. It is the main cause of increased pressure in the colon. The excess pressure causes the weak spots in the colon to bulge out and become diverticula.

Diverticulitis occurs when diverticula become infected or inflamed. Doctors are not certain what causes the infection. It may begin when stool or bacteria are caught in the diverticula. An attack of diverticulitis can develop suddenly and without warning.

What Are The Symptoms?

Diverticulosis

Most people with diverticulosis do not have any discomfort or symptoms. However, symptoms may include mild cramps, bloating and constipation. Other diseases such as irritable bowel syndrome (IBS) and stomach ulcers cause similar problems, so these symptoms do not always mean a person has diverticulosis. You should visit your doctor if you have any of these troubling symptoms.

Diverticulitis

The most common symptom of diverticulitis is abdominal pain. The pain is usually located around the left side of the lower abdomen. If infection is the cause, fever, nausea, vomiting, chills, cramping, diarrhea or constipation may occur as well. The severity of symptoms depends on the extent of the infection and any possible complications.

Are There Complications of Diverticulitis?

Diverticulitis can lead to complications such as infections, perforations or tears, blockages, or bleeding. These complications always require treatment to prevent them from progressing and causing more serious illness

Bleeding

Bleeding from diverticular disease is a rare complication. When diverticula bleed, blood may appear in the toilet or in the stool. Bleeding can be severe, but it often stops by itself and does not require any specific treatment. Doctors believe bleeding diverticula are caused by a small blood vessel in a diverticulum that weakens and finally bursts. If you have bleeding from the rectum, you should see your doctor. If the bleeding does not stop, surgery may be necessary.

Abscess, Perforation, and Peritonitis

The infection causing diverticulitis often clears up after a few days of treatment with antibiotics. If the condition gets worse, an abscess may form in the colon.

An abscess is an infected area with pus that may cause swelling and destroy tissue. Sometimes, the infected diverticula may develop small holes, called perforations. These perforations allow pus to leak out of the colon into the abdominal area. If the abscess is small and remains in the colon, it may clear up after treatment with antibiotics. If the abscess does not clear up with antibiotics, the doctor may need to drain it.

A large abscess can become a serious problem if the infection leaks out and contaminates areas outside the colon. Infection that spreads into the abdominal cavity is called peritonitis. Peritonitis requires immediate surgery to clean the abdominal cavity and remove the damaged part of the colon. Without surgery, peritonitis can be fatal.

Fistula

A fistula is an abnormal connection of tissue between two organs or between an organ and the skin. When damaged tissues come into contact with each other during infection, they sometimes stick together. If they heal that way, a fistula forms. When diverticulitis related infection spreads outside the colon, the colon's tissue may stick to nearby tissues. The most common organs involved are the urinary bladder, small intestine, gallbladder and skin.

The most common type of fistula occurs between the bladder and the colon. This type of fistula can result in a severe, long-lasting infection of the urinary tract. The problem can be corrected with surgery to remove the fistula and the affected part of the colon.

Intestinal Obstruction

The scarring caused by infection may cause partial or total blockage of the large intestine. When this happens, the colon is unable to move bowel contents along normally. When the obstruction totally blocks the intestine, emergency surgery is necessary. Partial blockage is not an emergency, so the surgery to correct it can be planned.

How Does the Doctor Diagnose Diverticular Disease?

To diagnose diverticular disease, the doctor asks about medical history, does a physical exam, and may perform one or more diagnostic tests, such as a colonoscopy or barium enema. Because most people do not have symptoms, diverticulosis is often found through tests ordered for another problem.

What Is the Treatment for Diverticular Disease?

A high-fiber diet and, occasionally, mild pain medications will help relieve symptoms in most cases. Sometimes an attack of diverticulitis is serious enough to require a hospital stay and possibly surgery.

Diverticulosis

Increasing the amount of fiber in the diet may reduce symptoms of diverticulosis and prevent complications such as diverticulitis. Fiber keeps stool soft and lowers pressure inside the colon so that bowel contents can move through easily. The American Dietetic Association recommends 20 to 35 grams of fiber each day.

Your doctor may also recommend taking a fiber supplement such as Citrucel or Metamucil once a day. These products provide about 4 to 6 grams of fiber in each dose.

Diverticulitis

Treatment for diverticulitis focuses on clearing up the infection and inflammation, resting the colon, and preventing or minimizing complications. An attack of diverticulitis without complications may respond to antibiotics within a few days if treated early.

To help the colon rest, your doctor may recommend bed rest and a liquid diet, along with a pain reliever or a drug to control muscle spasms in the colon.

An acute attack with severe pain or severe infection may require a hospital stay. Most acute cases of diverticulitis are treated with antibiotics and a liquid diet. The antibiotics are given by injection into a vein. In some cases, however, surgery may be necessary.